

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004470 A1

(51) International Patent Classification⁷: **C07D 207/34,**
A61K 31/40

Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **KRZYZANIAK, Joseph, Francis** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US).

(21) International Application Number: **PCT/IB02/01796**

(22) International Filing Date: **21 May 2002 (21.05.2002)**

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 201 Tabor Road, Morris Plains, NJ 07950 (US).

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
60/302,049 **29 June 2001 (29.06.2001)** **US**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*):
WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BYRN, Stephen, Robert** [US/US]; 824 Barlow Street, West Lafayette, IN 47906 (US). **COATES, David, Andrew** [US/US]; 2435 Neil Armstrong Dr. Apt 24, West Lafayette, IN 47906 (US). **GUSHURST, Karen, Sue** [US/US]; 807 Columbia Street, Lafayette, IN 47901 (US). **MORRISON, Henry, Grant, II** [US/US]; 2304 Osage Ct., Apt. 70, Lafayette, IN 47909 (US). **PARK, Aeri** [KR/US]; 1717 Southworth Branch Dr., West Lafayette, IN 47906 (US). **VLAHOVA, Petinka Ivanova** [BG/US]; 4702 Admirals Pointe Dr., Lafayette, IN 47909 (US). **LI, Zheng, Jane** [US/US];

Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/004470 A1

(54) Title: **CRYSTALLINE FORMS OF 'R-(R*,R*)-2-(4-FLUOROPHENYL)-BETA, DELTA-DIHYDROXY-5-(1-METHYLETHYL)-3-PHENYL-4-'PHENYLAMINO)CARBONYL-1H-PYRROLE-1-HEPTANOIC ACID CALCIUM SALT (2:1) (ATORVASTATIN)**

(57) Abstract: Novel crystalline forms of [R-(R*,R*)]-2-(4-fluorophenyl)-β, Δ;-dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid hemi calcium salt atorvastatin designated Form V, Form VI, Form VII, Form VIII, Form IX, Form X, Form XI, Form XII, Form XIII, Form XIV, Form XV, Form XVI, Form XVII, Form XVIII, and Form XIX are characterized by their X-ray powder diffraction, solid-state NMR, and/or Raman spectroscopy are described, as well as methods for the preparation and pharmaceutical composition of the same, which are useful as agents for treating hyperlipidemia, hypercholesterolemia, osteoporosis, and Alzheimer's disease.

EXPRESS MAIL NO. EV545395338US
DOCKET NO. PC25264A

CRYSTALLINE FORMS OF $R-(R^*, R^*)-2-(4\text{-FLUOROPHENYL})-\beta, \delta$ -DIHYDROXY-5-(1-METHYLETHYL)-3-PHENYL-4-[(PHENYLAMINO)CARBONYL]-1H-PYRROLE-1-HEPTANOIC ACID CALCIUM SALT (2:1) (ATORVASTATIN)

5

FIELD OF THE INVENTION

The present invention relates to novel crystalline forms of atorvastatin which is known by the chemical name $R-(R^*, R^*)-2-(4\text{-fluorophenyl})-\beta, \delta$ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid hemi calcium salt useful as pharmaceutical agents, to methods for their production and isolation, to pharmaceutical compositions which include these compounds and a pharmaceutically acceptable carrier, as well as methods of using such compositions to treat subjects, including human subjects, suffering from hyperlipidemia, hypercholesterolemia, osteoporosis, and Alzheimer's disease.

15

BACKGROUND OF THE INVENTION


The conversion of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) to mevalonate is an early and rate-limiting step in the cholesterol biosynthetic pathway. This step is catalyzed by the enzyme HMG-CoA reductase. Statins inhibit HMG-CoA reductase from catalyzing this conversion. As such, statins are collectively potent lipid lowering agents.

20

Atorvastatin calcium, disclosed in United States Patent No. 5,273,995, which is incorporated herein by reference, is currently sold as Lipitor® having the chemical name $R-(R^*, R^*)-2-(4\text{-fluorophenyl})-\beta, \delta$ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid calcium salt (2:1) trihydrate and the formula

25

-2-



Atorvastatin calcium is a selective, competitive inhibitor of HMG-CoA reductase. As such, atorvastatin calcium is a potent lipid lowering compound and is thus useful as a hypolipidemic and/or hypocholesterolemic agent.

5 United States Patent Number 4,681,893, which is incorporated herein by reference, discloses certain *trans*-6-[2-(3- or 4-carboxamido-substituted-pyrrol-1-yl)alkyl]-4-hydroxy-pyran-2-ones including *trans* (\pm)-5-(4-fluorophenyl)-2-(1-methylethyl)-N, 4-diphenyl-1-[(2-tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide.

10 United States Patent Number 5,273,995, which is herein incorporated by reference, discloses the enantiomer having the R form of the ring-opened acid of *trans*-5-(4-fluorophenyl)-2-(1-methylethyl)-N, 4-diphenyl-1-[(2-tetrahydro-4-hydroxy-6-oxo-2H-pyran-2-yl)ethyl]-1H-pyrrole-3-carboxamide, ie, [R-(R*,R*)]-2-(4-fluorophenyl)- β , δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)-carbonyl]-1H-pyrrole-1-heptanoic acid which is atorvastatin.

15 United States Patent Numbers 5,003,080; 5,097,045; 5,103,024; 5,124,482; 5,149,837; 5,155,251; 5,216,174; 5,245,047; 5,248,793; 5,280,126; 5,397,792; 5,342,952; 5,298,627; 5,446,054; 5,470,981; 5,489,690; 5,489,691; 5,510,488; 5,998,633; and 6,087,511, which are herein incorporated by reference, disclose various processes and key intermediates for preparing amorphous atorvastatin. Amorphous atorvastatin has unsuitable filtration and drying characteristics for large-scale production and must be protected from heat, light, oxygen, and moisture.

20

-3-

Crystalline forms of atorvastatin calcium are disclosed in United States Patent Numbers 5,969,156 and 6,121,461 which are herein incorporated by reference.

International Published Patent Application Number WO 01/36384
5 allegedly discloses a polymorphic form of atorvastatin calcium.

Stable oral formulations of atorvastatin calcium are disclosed in United States Patent Numbers 5,686,104 and 6,126,971.

Atorvastatin is prepared as its calcium salt, ie, [R-(R*,R*)]-2-(4-fluorophenyl)- β , δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-
10 [(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid calcium salt (2:1). The calcium salt is desirable since it enables atorvastatin to be conveniently formulated in, for example, tablets, capsules, lozenges, powders, and the like for oral administration. Additionally, there is a need to produce atorvastatin in a pure and crystalline form to enable formulations to meet exacting pharmaceutical
15 requirements and specifications.

Furthermore, the process by which atorvastatin is produced needs to be one which is amenable to large-scale production. Additionally, it is desirable that the product should be in a form that is readily filterable and easily dried. Finally, it is economically desirable that the product be stable for extended periods of time
20 without the need for specialized storage conditions.

We have now surprisingly and unexpectedly found novel crystalline forms of atorvastatin. Thus, the present invention provides atorvastatin in new crystalline forms designated Forms V, VI, VII, VIII, IX, X, XI, XII, XIII, XIV, XV, XVI, XVII, XVIII, and XIX. The new crystalline forms of atorvastatin are purer, more
25 stable, or have advantageous manufacturing properties than the amorphous product.

SUMMARY OF THE INVENTION

Accordingly, the present invention is directed to crystalline Form V atorvastatin and hydrates thereof characterized by the following X-ray powder
30 diffraction pattern expressed in terms of the 2 θ and relative intensities with a

-4-

relative intensity of >10% measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

| 2 θ | Relative Intensity (>10%) ^a |
|-------------|---|
| 4.9 (broad) | 9 |
| 6.0 | 15 |
| 7.0 | 100 |
| 8.0 (broad) | 20 |
| 8.6 | 57 |
| 9.9 | 22 |
| 16.6 | 42 |
| 19.0 | 27 |
| 21.1 | 35 |

^a Relative intensity of 4.9 (broad) 2 θ is 9.

Additionally, the following X-ray powder diffraction pattern of crystalline Form V atorvastatin expressed in terms of the 2 θ values was measured on an Inel (capillary) diffractometer:

5

| 2 θ |
|-------------|
| 5.0 |
| 6.1 |
| 7.5 |
| 8.4 (broad) |
| 8.7 (broad) |
| 9.9 |
| 16.7 |
| 19.0 |
| 21.2 |

-5-

Further, the present invention is directed to crystalline Form V atorvastatin and hydrates thereof characterized by the following solid-state ^{13}C nuclear magnetic resonance (ssNMR) spectrum wherein chemical shift is expressed in parts per million:

| Assignment | Chemical Shift |
|-------------------------------------|----------------|
| C12 or C25 | 185.7 |
| C12 or C25 | 176.8 |
| C16 | 166.9 |
| <u>Aromatic Carbons</u> | 138.7 |
| C2-C5, C13-C18, C19-C24, C27-C32 | 136.3 |
| | 133.0 |
| | 128.4 |
| | 122.0 |
| | 117.0 |
| | 116.3 |
| C8, C10 | 68.0 |
| <u>Methylene Carbons</u> | 43.1 |
| C6, C7, C9, C11 | |
| C33 | 25.6 |
| C34 | 19.9 |

- 5 Additionally, the present invention is directed to crystalline Form V atorvastatin and hydrates thereof characterized by the following Raman spectrum having peaks expressed in cm^{-1} :

-6-

3062

1652

1604

1528

1478

1440

1413

1397

1368

1158

1034

1001

825

245

224

130

In a preferred embodiment of the first aspect of the invention, crystalline Form V atorvastatin is a trihydrate.

In a second aspect, the present invention is directed to crystalline Form VI atorvastatin and hydrates thereof characterized by the following X-ray powder
5 diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of >10% measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

-7-

| 2 θ | Relative Intensity (>10%) ^a |
|--------------|---|
| 7.2 | 11 |
| 8.3 | 77 |
| 11.0 | 20 |
| 12.4 | 11 |
| 13.8 | 9 |
| 16.8 | 14 |
| 18.5 | 100 |
| 19.7 (broad) | 22 |
| 20.9 | 14 |
| 25.0 (broad) | 15 |

^a Relative intensity of 13.8 (broad) 2 θ is 9.

Additionally, the following X-ray powder diffraction pattern of crystalline Form VI atorvastatin expressed in terms of the 2 θ values was measured on an Inel (capillary) diffractometer:

| 2 θ |
|--------------|
| 7.3 |
| 8.5 |
| 11.2 |
| 12.7 |
| 14.0 |
| 17.1 (broad) |
| 18.7 |
| 19.9 |
| 21.1 (broad) |
| 25.2 (broad) |

Further, the present invention is directed to crystalline Form VI
 5 atorvastatin and hydrates thereof characterized by the following solid-state

-8-

^{13}C nuclear magnetic resonance spectrum wherein chemical shift is expressed in parts per million:

| Assignment | Chemical Shift |
|-------------------------------------|----------------|
| C12 or C25 | 176.5 |
| C16 or C12 or C25 | 168.2 |
| C16 or C12 or C25 | 163.1 |
| C16 or C12 or C25 | 159.8 |
| <u>Aromatic Carbons</u> | 136.8 |
| C2-C5, C13-C18, C19-C24, C27-C32 | 127.8 |
| | 122.3 |
| | 118.8 |
| | 113.7 |
| C8, C10 | 88.2 |
| C8, C10 | 79.3 |
| | 70.5 |
| <u>Methylene Carbons</u> | 43.3 |
| C6, C7, C9, C11 | 36.9 |
| | 31.9 |
| C33, C34 | 25.9 |
| C33, C34 | 22.5 |

In a third aspect, the present invention is directed to crystalline Form VII atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of >10% measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

-9-

| 2 θ | Relative Intensity (>10%) |
|--------------|------------------------------|
| 8.6 | 76 |
| 10.2 | 70 |
| 12.4 (broad) | 12 |
| 12.8 (broad) | 15 |
| 17.6 | 20 |
| 18.3 (broad) | 43 |
| 19.3 | 100 |
| 22.2 (broad) | 14 |
| 23.4 (broad) | 23 |
| 23.8 (broad) | 26 |
| 25.5 (broad) | 16 |

Additionally, the following X-ray powder diffraction pattern of crystalline Form VII atorvastatin expressed in terms of the 2 θ values was measured on an Inel (capillary) diffractometer:

| 2 θ |
|------------|
| 8.7 |
| 10.2 |
| 12.4 |
| 12.9 |
| 17.6 |
| 18.4 |
| 19.4 |
| 22.2 |
| 23.5 |
| 23.9 |
| 25.6 |

-10-

Further, the present invention is directed to crystalline Form VII atorvastatin and hydrates thereof characterized by the following solid-state ^{13}C nuclear magnetic resonance spectrum wherein chemical shift is expressed in parts per million:

| Assignment | Chemical Shift |
|-------------------------------------|----------------|
| C12 or C25 | 186.5 |
| C12 or C25 | 183.3 |
| C12 or C25 | 176.8 |
| C16 | 166.5 |
| | 159.2 |
| <u>Aromatic Carbons</u> | 137.6 |
| C2-C5, C13-C18, C19-C24, C27-C32 | 128.3 |
| | 122.3 |
| | 119.2 |
| C8, C10 | 74.5 |
| C8, C10 | 70.3 |
| C8, C10 | 68.3 |
| C8, C10 | 66.2 |
| <u>Methylene Carbons</u> | 43.5 |
| C6, C7, C9, C11 | 40.3 |
| C33, C34 | 26.3 |
| C33, C34 | 24.9 |
| C33, C34 | 20.2 |

5

Additionally, the present invention is directed to crystalline Form VII atorvastatin and hydrates thereof characterized by the following Raman spectrum having peaks expressed in cm^{-1} :

-11-

Raman Spectrum

3060

2927

1649

1603

1524

1476

1412

1397

1368

1159

1034

998

824

114

In a preferred embodiment of the third aspect of the invention, crystalline Form VII atorvastatin is a sesquihydrate.

In a fourth aspect, the present invention is directed to crystalline Form VIII atorvastatin and hydrates thereof characterized by the following X-ray powder
5 diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of $>10\%$ measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

-12-

| 2 θ | Relative Intensity (>10%) ^a |
|--------------|---|
| 7.5 | 61 |
| 9.2 | 29 |
| 10.0 | 16 |
| 12.1 | 10 |
| 12.8 | 6 |
| 13.8 | 4 |
| 15.1 | 13 |
| 16.7 (broad) | 64 |
| 18.6 (broad) | 100 |
| 20.3 (broad) | 79 |
| 21.2 | 24 |
| 21.9 | 30 |
| 22.4 | 19 |
| 25.8 | 33 |
| 26.5 | 20 |
| 27.4 (broad) | 38 |
| 30.5 | 20 |

^a Relative intensity of 12.8 2 θ is 6 and
13.8 2 θ is 4.

Additionally, the following X-ray powder diffraction pattern of crystalline Form VIII atorvastatin expressed in terms of the 2 θ values was measured on an Inel (capillary) diffractometer:

| 2 θ |
|------------|
| 7.5 |
| 9.3 |
| 10.1 |
| 12.2 |
| 12.8 |

| -13- |
|--------------|
| 2θ |
| 13.8 |
| 15.1 |
| 16.6-16.9 |
| 18.5-18.9 |
| 20.2-20.6 |
| 21.3 |
| 22.0 |
| 22.5 |
| 25.9 |
| 26.5 |
| 27.4 (broad) |
| 30.6 |

Further, the present invention is directed to crystalline Form VIII atorvastatin and hydrates thereof characterized by the following solid-state ^{13}C nuclear magnetic resonance spectrum wherein chemical shift is expressed in parts per million:

| Assignment | Chemical Shift |
|-------------------------------------|----------------|
| C12 or C25 | 186.1 |
| C12 or C25 | 179.5 |
| C16 | 167.9 |
| C16 | 161.0 |
| <u>Aromatic Carbons</u> | 139.4 |
| C2-C5, C13-C18, C19-C24, C27-C32 | 132.9 |
| | 128.7 |
| | 124.7 |
| | 121.8 |
| | 116.6 |
| C8, C10 | 67.0 |

-14-

| Assignment | Chemical Shift |
|--------------------------|----------------|
| <u>Methylene Carbons</u> | 43.3 |
| C6, C7, C9, C11 | |
| C33, C34 | 26.7 |
| C33, C34 | 24.7 |
| C33, C34 | 20.9 |
| C33, C34 | 20.1 |

Additionally, the present invention is directed to crystalline Form VIII atorvastatin and hydrates thereof characterized by the following Raman spectrum having peaks expressed in cm^{-1} :

| Raman Spectrum |
|----------------|
| 3065 |
| 2923 |
| 1658 |
| 1603 |
| 1531 |
| 1510 |
| 1481 |
| 1413 |
| 997 |
| 121 |

5 In a preferred embodiment of the fourth aspect of the invention, crystalline Form VIII atorvastatin is a dihydrate.

In a fifth aspect, the present invention is directed to crystalline Form IX atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of $>10\%$ measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

10

-15-

| 2 θ | Relative Intensity (>10%) |
|-------------------|------------------------------|
| 8.8 | 50 |
| 9.4 (broad) | 32 |
| 11.2-11.7 (broad) | 26 |
| 16.7 | 59 |
| 17.5 (broad) | 33 |
| 19.3 (broad) | 55 |
| 21.4 (broad) | 100 |
| 22.4 (broad) | 33 |
| 23.2 (broad) | 63 |
| 29.0 (broad) | 15 |

Additionally, the following X-ray powder diffraction pattern of crystalline Form IX atorvastatin expressed in terms of the 2 θ values was measured on an Inel (capillary) diffractometer:

| 2 θ |
|---------------------|
| 9.0 |
| 9.4 |
| 10.0 - 10.5 (broad) |
| 11.8 - 12.0 (broad) |
| 16.9 |
| 17.5 (broad) |
| 19.4 (broad) |
| 21.6 (broad) |
| 22.6 (broad) |
| 23.2 (broad) |
| 29.4 (broad) |

In a sixth aspect, the present invention is directed to crystalline Form X
 5 atorvastatin and hydrates thereof characterized by the following X-ray powder

-16-

diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of >10% measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

| 2θ | Relative Intensity (>10%) |
|--------------|------------------------------|
| 4.7 | 35 |
| 5.2 | 24 |
| 5.8 | 11 |
| 6.9 | 13 |
| 7.9 | 53 |
| 9.2 | 56 |
| 9.5 | 50 |
| 10.3 (broad) | 13 |
| 11.8 | 20 |
| 16.1 | 13 |
| 16.9 | 39 |
| 19.1 | 100 |
| 19.8 | 71 |
| 21.4 | 49 |
| 22.3 (broad) | 36 |
| 23.7 (broad) | 37 |
| 24.4 | 15 |
| 28.7 | 31 |

5 Additionally, the following X-ray powder diffraction pattern of crystalline Form X atorvastatin expressed in terms of the 2θ values was measured on an Inel (capillary) diffractometer:

| |
|-------------------|
| -17- |
| 20 |
| 4.7 |
| 5.2 |
| 5.8 |
| 6.9 |
| 7.9 |
| 9.2 |
| 9.6 |
| 10.2-10.4 |
| 11.9 |
| 16.2 |
| 16.9 |
| 19.1 |
| 19.9 |
| 21.5 |
| 22.3-22.6 |
| 23.7-24.0 (broad) |
| 24.5 |
| 28.8 |

Further, the present invention is directed to crystalline Form X atorvastatin and hydrates thereof characterized by the following solid-state ^{13}C nuclear magnetic resonance spectrum wherein chemical shift is expressed in parts per million:

| Assignment | Chemical Shift |
|------------|----------------|
| C12 or C25 | 187.0 |
| C12 or C25 | 179.5 |
| C16 | 165.5 |
| C16 | 159.4 |

-18-

| Assignment | Chemical Shift |
|--------------------------|----------------|
| <u>Aromatic Carbons</u> | 137.9 |
| C2-C5, C13-C18, | 134.8 |
| C19-C24, C27-C32 | 129.4 |
| | 127.9 |
| | 123.2 |
| | 119.9 |
| C8, C10 | 71.1 |
| <u>Methylene Carbons</u> | 43.7 |
| C6, C7, C9, C11 | 40.9 |
| C33 | 26.4 |
| | 25.3 |
| C34 | 20.3 |
| | 18.3 |

Additionally, the present invention is directed crystalline Form X atorvastatin and hydrates thereof characterized by the following Raman spectrum having peaks expressed in cm^{-1} :

| Raman Spectrum |
|----------------|
| 3062 |
| 2911 |
| 1650 |
| 1603 |
| 1525 |
| 1478 |
| 1411 |
| 1369 |
| 1240 |
| 1158 |
| 1034 |
| 999 |

-19-

Raman Spectrum

824

116

In a preferred embodiment of the sixth aspect of the invention, crystalline Form X atorvastatin is a trihydrate.

In a seventh aspect, the present invention is directed to crystalline Form XI atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of >10% measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

| 2θ | Relative Intensity (>10%) |
|-------------------|------------------------------|
| 10.8 (broad) | 58 |
| 12.0 | 12 |
| 13.5 | 11 |
| 16.5 | 52 |
| 17.6-18.0 (broad) | 35 |
| 19.7 | 82 |
| 22.3 | 100 |
| 23.2 | 26 |
| 24.4 | 28 |
| 25.8 | 17 |
| 26.5 | 30 |
| 27.3 | 31 |
| 28.7 | 19 |
| 29.5 | 12 |
| 30.9 (broad) | 17 |
| 32.8 (broad) | 11 |
| 33.6 (broad) | 15 |
| 36.0 (broad) | 15 |
| 38.5 (broad) | 14 |

-20-

In an eighth aspect, the present invention is directed to crystalline Form XII atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of $>10\%$ measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

5

| 2θ | Relative Intensity ($>10\%$) ^a |
|-------------------|--|
| 5.4 | 11 |
| 7.7 | 24 |
| 8.0 | 25 |
| 8.6 | 42 |
| 8.9 | 25 |
| 9.9 | 36 |
| 10.4 (broad) | 24 |
| 12.5 | 18 |
| 13.9 (broad) | 9 |
| 16.2 | 10 |
| 17.8 | 70 |
| 19.4 | 100 |
| 20.8 | 51 |
| 21.7 | 13 |
| 22.4-22.6 (broad) | 18 |
| 24.3 | 19 |
| 25.5 | 24 |
| 26.2 | 11 |
| 27.1 | 8 |

^a Relative intensity of 13.9 (broad) 2θ is 9 and 27.1 2θ is 8.

Additionally, the following X-ray powder diffraction pattern of crystalline Form XII atorvastatin expressed in terms of the 2θ values was measured on an Inel (capillary) diffractometer:

| |
|---------------------|
| -21- |
| 2 θ |
| 5.4 |
| 7.7 |
| 8.1 |
| 8.6 |
| 8.9 |
| 10.0 |
| 10.5 |
| 12.6 |
| 14.0 (broad) |
| 16.2 |
| 17.9 |
| 19.4 |
| 20.9 |
| 21.8 |
| 22.5 - 22.8 (broad) |
| 24.4 |
| 25.6 |
| 26.4 |
| 27.2 |

Additionally, the present invention is directed crystalline Form XII atorvastatin and hydrates thereof characterized by the following Raman spectrum having peaks expressed in cm^{-1} :

| |
|----------------|
| Raman Spectrum |
| 3064 |
| 2973 |
| 2926 |
| 1652 |
| 1603 |
| 1527 |

-22-

| Raman Spectrum |
|----------------|
| 1470 |
| 1410 |
| 1367 |
| 1240 |
| 1159 |
| 1034 |
| 1002 |
| 823 |

In a ninth aspect, the present invention is directed to crystalline Form XIII atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of >10% measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

5

| 2θ | Relative Intensity (>10%) |
|--------------|------------------------------|
| 8.4 | 100 |
| 8.9 | 82 |
| 15.7 (broad) | 45 |
| 16.4 (broad) | 46 |
| 17.6 (broad) | 57 |
| 18.1 (broad) | 62 |
| 19.7 (broad) | 58 |
| 20.8 (broad) | 91 |
| 23.8 (broad) | 57 |

In a tenth aspect, the present invention is directed to crystalline Form XIV atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a

-23-

relative intensity of >10% measured on a Bruker D5000 diffractometer with
CuK α radiation:

| 2 θ | Relative Intensity (>10%) |
|---------------------|------------------------------|
| 5.4 | 41 |
| 6.7 | 31 |
| 7.7 | 100 |
| 8.1 | 35 |
| 9.0 | 65 |
| 16.5 (broad) | 15 |
| 17.6 (broad) | 17 |
| 18.0 – 18.7 (broad) | 21 |
| 19.5 (broad) | 18 |

In an eleventh aspect, the present invention is directed to crystalline
Form XV atorvastatin and hydrates thereof characterized by the following X-ray
5 powder diffraction pattern expressed in terms of the 2 θ and relative intensities
with a relative intensity of >10% measured on a Bruker D5000 diffractometer
with CuK α radiation:

| 2 θ | Relative Intensity (>10%) |
|---------------------|------------------------------|
| 5.7 | 26 |
| 6.1 | 21 |
| 6.8 | 18 |
| 7.5 | 39 |
| 8.1 | 39 |
| 8.5 | 42 |
| 9.5 | 33 |
| 10.5 (broad) | 18 |
| 19.1 - 19.6 (broad) | 32 |

-24-

In a twelfth aspect, the present invention is directed to crystalline Form XVI atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of >10% measured on a Bruker D5000 diffractometer with $\text{CuK}\alpha$ radiation:

5

| 2θ | Relative Intensity (>10%) |
|---------------------|------------------------------|
| 5.2 | 37 |
| 6.4 | 34 |
| 7.5 | 100 |
| 8.7 | 79 |
| 10.5 (broad) | 19 |
| 12.0 (broad) | 10 |
| 12.7 (broad) | 17 |
| 16.7 | 26 |
| 18.3 (broad) | 27 |
| 19.5 | 23 |
| 20.1 - 20.4 (broad) | 37 |
| 21.2 - 21.9 (broad) | 32 |
| 22.9 - 23.3 (broad) | 38 |
| 24.4 - 25.0 (broad) | 35 |

Additionally, the following X-ray powder diffraction pattern of crystalline Form XVI atorvastatin expressed in terms of the 2θ values was measured on a Shimadzu diffractometer with $\text{CuK}\alpha$ radiation:

| 2θ |
|-----------|
| 7.6 |
| 8.8 |
| 10.2 |
| 12.5 |

| -25- |
|-----------|
| 2θ |
| 16.8 |
| 18.2 |
| 19.3 |
| 20.5 |
| 23.0 |
| 24.8 |

In addition, the following X-ray powder diffraction pattern of crystalline Form XVI atorvastatin expressed in terms of the 2θ values was measured on an Inel (capillary) diffractometer:

| 2θ |
|---------------------|
| 5.1 |
| 6.2 |
| 7.3 |
| 8.7 |
| 10.2 (broad) |
| 12.0 (broad) |
| 12.7 (broad) |
| 16.7 |
| 18.0 (broad) |
| 19.5 (broad) |
| 20.0 - 20.5 (broad) |
| 21.5 - 21.6 (broad) |
| 22.9 - 23.3 (broad) |
| 24.0 - 25.0 (broad) |

5 In a thirteenth aspect, the present invention is directed to crystalline Form XVII atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities

-26-

with a relative intensity of >10% measured on a Bruker D5000 diffractometer with $\text{CuK}\alpha$ radiation:

| 2 θ | Relative Intensity (>10%) |
|---------------------|------------------------------|
| 5.0 | 27 |
| 6.1 | 33 |
| 7.3 | 100 |
| 7.9 | 30 |
| 8.5 | 29 |
| 9.1 | 22 |
| 10.0 | 45 |
| 12.1 (broad) | 24 |
| 14.8 | 17 |
| 16.0 - 16.5 (broad) | 20 |
| 17.5 (broad) | 28 |
| 19.0 (broad) | 46 |
| 19.5 | 65 |
| 20.2 (broad) | 47 |
| 21.3 | 64 |
| 21.6 | 55 |
| 22.0 | 45 |

5 In a fourteenth aspect, the present invention is directed to crystalline Form XVIII atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2 θ and relative intensities with a relative intensity of >10% measured on a Bruker D5000 diffractometer with $\text{CuK}\alpha$ radiation:

-27-

| 2 θ | Relative Intensity (>10%) |
|--------------|------------------------------|
| 8.0 | 100 |
| 9.2 (broad) | 52 |
| 9.7 (broad) | 40 |
| 12.1 | 24 |
| 16.6 (broad) | 48 |
| 18.5 | 67 |

Additionally, the following X-ray powder diffraction pattern of crystalline Form XVIII atorvastatin expressed in terms of the 2 θ values was measured on a Shimadzu diffractometer with CuK α radiation:

| 2 θ |
|------------|
| 7.7 |
| 9.3 |
| 9.9 |
| 12.2 |
| 16.8 |
| 18.5 |

5 In addition, the following X-ray powder diffraction pattern of crystalline Form XVIII atorvastatin expressed in terms of the 2 θ values was measured on an Inel (capillary) diffractometer:

| 2 θ |
|--------------|
| 7.9 |
| 9.2 (broad) |
| 9.8 (broad) |
| 12.2 (broad) |
| 16.7 (broad) |
| 18.5 |

-28-

In a fifteenth aspect, the present invention is directed to crystalline Form XIX atorvastatin and hydrates thereof characterized by the following X-ray powder diffraction pattern expressed in terms of the 2θ and relative intensities with a relative intensity of $>10\%$ measured on a Bruker D5000 diffractometer with $\text{CuK}\alpha$ radiation:

| 2θ | Relative Intensity ($>10\%$) |
|--------------|-----------------------------------|
| 5.2 | 32 |
| 6.3 | 28 |
| 7.0 | 100 |
| 8.6 | 74 |
| 10.5 | 34 |
| 11.6 (broad) | 26 |
| 12.7 (broad) | 35 |
| 14.0 | 15 |
| 16.7 (broad) | 30 |
| 18.9 | 86 |
| 20.8 | 94 |
| 23.6 (broad) | 38 |
| 25.5 (broad) | 32 |

As inhibitors of HMG-CoA reductase, the novel crystalline forms of atorvastatin are useful hypolipidemic and hypocholesterolemic agents as well as agents in the treatment of osteoporosis and Alzheimer's disease.

A still further embodiment of the present invention is a pharmaceutical composition for administering an effective amount of crystalline Form V, Form VI, Form VII, Form VIII, Form IX, Form X, Form XI, Form XII, Form XIII, Form XIV, Form XV, Form XVI, Form XVII, Form XVIII, or Form XIX atorvastatin in unit dosage form in the treatment methods mentioned above. Finally, the present invention is directed to methods for production of Form V, Form VI, Form VII, Form VIII, Form IX, Form X, Form XI, Form XII,

-29-

Form XIII, Form XIV, Form XV, Form XVI, Form XVII, Form XVIII, or
Form XIX atorvastatin.

BRIEF DESCRIPTION OF THE DRAWINGS

5 The invention is further described by the following nonlimiting examples
which refer to the accompanying Figures 1 to 35, short particulars of which are
given below.

Figure 1

Diffractiongram of Form V atorvastatin carried out on Shimadzu XRD-6000
diffractometer.

10 Figure 2

Diffractiongram of Form VI atorvastatin carried out on Shimadzu
XRD-6000 diffractometer.

Figure 3

15 Diffractiongram of Form VII atorvastatin carried out on Shimadzu
XRD-6000 diffractometer.

Figure 4

Diffractiongram of Form VIII atorvastatin carried out on Shimadzu
XRD-6000 diffractometer.

Figure 5

20 Diffractiongram of Form IX atorvastatin carried out on Shimadzu
XRD-6000 diffractometer.

Figure 6

Diffractiongram of Form X atorvastatin carried out on Shimadzu XRD-6000
diffractometer.

25 Figure 7

Diffractiongram of Form XI atorvastatin carried out on Shimadzu
XRD-6000 diffractometer.

-30-

Figure 8

Diffractiongram of Form XII atorvastatin carried out on Shimadzu XRD-6000 diffractometer.

Figure 9

5 Diffractiongram of Form XIII atorvastatin carried out on Shimadzu XRD-6000 diffractometer.

Figure 10

Diffractiongram of Form XIV atorvastatin carried out on Bruker D 5000 diffractometer.

10 Figure 11

Diffractiongram of Form XV atorvastatin carried out on Bruker D 5000 diffractometer.

Figure 12

15 Diffractiongram of Form XVI atorvastatin carried out on Bruker D 5000 diffractometer.

Figure 13

Diffractiongram of Form XVII atorvastatin carried out on Bruker D 5000 diffractometer.

Figure 14

20 Diffractiongram of Form XVIII atorvastatin carried out on Bruker D 5000 diffractometer.

Figure 15

Diffractiongram of Form XIX atorvastatin carried out on Bruker D 5000 diffractometer.

25 Figure 16

Diffractiongram of Form V atorvastatin carried out on Inel XRG-3000 diffractometer.

-31-

Figure 17

Diffractiongram of Form VI atorvastatin carried out on Inel XRG-3000 diffractometer.

Figure 18

5 Diffractiongram of Form VII atorvastatin carried out on Inel XRG-3000 diffractometer.

Figure 19

Diffractiongram of Form VIII atorvastatin carried out on Inel XRG-3000 diffractometer.

10 Figure 20

Diffractiongram of Form IX atorvastatin carried out on Inel XRG-3000 diffractometer.

Figure 21

15 Diffractiongram of Form X atorvastatin carried out on Inel XRG-3000 diffractometer.

Figure 22

Diffractiongram of Form XII atorvastatin carried out on Inel XRG-3000 diffractometer.

Figure 23

20 Diffractiongram of Form XVI atorvastatin carried out on Inel XRG-3000 diffractometer.

Figure 24

Diffractiongram of Form XVIII atorvastatin carried out on Inel XRG-3000 diffractometer.

25 Figure 25

Solid-state ^{13}C nuclear magnetic resonance spectrum with spinning side bands identified by an asterisk of Form V atorvastatin.

-32-

Figure 26

Solid-state ^{13}C nuclear magnetic resonance spectrum with spinning side bands identified by an asterisk of Form VI atorvastatin.

Figure 27

5 Solid-state ^{13}C nuclear magnetic resonance spectrum with spinning side bands identified by an asterisk of Form VII atorvastatin.

Figure 28

Solid-state ^{13}C nuclear magnetic resonance spectrum with spinning side bands identified by an asterisk of Form VIII atorvastatin.

10 Figure 29

Solid-state ^{13}C nuclear magnetic resonance spectrum of Form X atorvastatin.

Figure 30

Raman spectrum of Form V.

15 Figure 31

Raman spectrum of Form VI.

Figure 32

Raman spectrum of Form VII.

Figure 33

20 Raman spectrum of Form VIII.

Figure 34

Raman spectrum of Form X.

Figure 35

Raman spectrum of Form XII.

25

DETAILED DESCRIPTION OF THE INVENTION

Crystalline Form V, Form VI, Form VII, Form VIII, Form IX, Form X, Form XI, Form XII, Form XIII, Form XIV, Form XV, Form XVI, Form XVII,

Form XVIII, and Form XIX atorvastatin may be characterized by their X-ray powder diffraction patterns, by their solid state nuclear magnetic resonance spectra (NMR), and/or their Raman spectra.

X-RAY POWDER DIFFRACTION

5 Forms V, VI, VII, VIII, IX, X, XI, XII, XIII, XIV, XV, XVI, XVII, XVIII, and XIX

Forms V, VI, VII, VIII, IX, X, XI, XII, XIII, XIV, XV, XVI, XVII, XVIII, or XIX atorvastatin were characterized by their X-ray powder diffraction pattern. Thus, the X-ray diffraction patterns of Forms V, VI, VII, VIII, IX, X, XI, XII, or
10 Form XIII atorvastatin were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using $\text{CuK}\alpha$ radiation. The instrument is equipped with a fine-focus X-ray tube. The tube voltage and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° , and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation
15 detector. A theta-two theta continuous scan at $3^\circ/\text{min}$ ($0.4 \text{ sec}/0.02^\circ$ step) from 2.5 to $40^\circ 2\theta$ was used. A silicon standard was analyzed each day to check the instrument alignment. The X-ray diffraction patterns of Forms XIV, XV, XVI, XVII, XVIII, and XIX were carried out on a Bruker D5000 diffractometer using copper radiation, fixed slits (1.0, 1.0, 0.6 mm), and a Kevex solid state detector.
20 Data was collected from 3.0 to 40.0 degrees in 2θ using a step size of 0.04 degrees and a step time of 1.0 seconds. It should be noted that Bruker Instruments purchased Siemens; thus, a Bruker D 5000 instrument is essentially the same as a Siemens D 5000.

25 The X-ray diffraction patterns of Forms V, VI, VII, VIII, IX, X, XII, XVI, and XVIII were also carried out on an Inel diffractometer. X-ray diffraction analyses were carried out on an Inel XRG-3000 diffractometer, equipped with a Curved Position Sensitive (CPS) detector with a 2θ range of 120 degrees. Real time data were collected using $\text{CuK}\alpha$ radiation starting at approximately $4^\circ 2\theta$ at a resolution of $0.03^\circ 2\theta$. The tube voltage and amperage were set to 40 kV and

-34-

30 mA, respectively. Samples were prepared for analysis by packing them into thin-walled glass capillaries. Each capillary was mounted onto a goniometer head that is motorized to permit spinning of the capillary during data acquisition. Instrument calibration was performed daily using a silicon reference standard. The Inel diffractograms for the available forms are shown in the figures without baseline subtraction. Calculating the intensities from these diffractograms is within the skill of the art and involves using baseline subtraction to account for background scattering (e.g., scattering from the capillary).

To perform an X-ray powder diffraction measurement on a Shimadzu or Bruker instrument like the ones used for measurements reported herein, the sample is typically placed into a holder which has a cavity. The sample powder is pressed by a glass slide or equivalent to ensure a random surface and proper sample height. The sample holder is then placed into the instrument (Shimadzu or Bruker). The source of the X-ray beam is positioned over the sample, initially at a small angle relative to the plane of the holder, and moved through an arc that continuously increases the angle between the incident beam and the plane of the holder. Measurement differences associated with such X-ray powder analyses result from a variety of factors including: (a) errors in sample preparation (e.g., sample height), (b) instrument errors (e.g., flat sample errors), (c) calibration errors, (d) operator errors (including those errors present when determining the peak locations), and (e) preferred orientation. Calibration errors and sample height errors often result in a shift of all the peaks in the same direction and by the same amount. Small differences in sample height on a flat holder lead to large displacements in XRPD peak positions. A systematic study showed that, using a Shimadzu XRD-6000 in the typical Bragg-Brentano configuration, sample height differences of 1 mm led to peak shifts as high as $1^\circ 2\theta$ (Chen, *et al.*, *J. Pharmaceutical and Biomedical Analysis*, 2001;26:63). These shifts can be identified from the X-ray diffractogram and can be eliminated by compensating for the shift (applying a systematic correction factor to all peak position values) or recalibrating the instrument. In contrast, the Inel instrument used herein places the sample in a capillary which is positioned at the center of the instrument. This minimizes sample height errors (a) and preferred orientation (e). Since, when

-35-

using capillaries, the sample height is not established manually, the peak locations from the Inel measurements are typically more accurate than those from the Shimadzu or the Bruker instrument. As mentioned above, it is possible to rectify measurements from the various machines by applying a systematic correction factor to bring the peak positions into agreement. In general, this correction factor will bring the peak positions from the Shimadzu and Bruker into agreement with the Inel and will be in the range of 0 to 0.2 °2 θ .

Table 1 lists the 2 θ and relative intensities of all lines in the sample with a relative intensity of >10% for crystalline Forms V-XIX atorvastatin. The numbers listed in this table are rounded numbers.

Table 1. Intensities and Peak Locations of All Diffraction Lines With Relative Intensity Greater Than 10%^a for Forms V to XIX
(Measured on Shimadzu Diffractometer)

(Page 1 of 3)

| Form V | | Form VI | | Form VII | | Form VIII | | Form IX | | Form X | | Form XI | | Form XII | |
|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|
| 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) |
| 4.9* | 9 | 7.2 | 11 | 8.6 | 76 | 7.5 | 61 | 8.8 | 50 | 4.7 | 35 | 10.8* | 58 | 5.4 | 11 |
| 6.0 | 15 | 8.3 | 77 | 10.2 | 70 | 9.2 | 29 | 9.4* | 32 | 5.2 | 24 | 12.0 | 12 | 7.7 | 24 |
| 7.0 | 100 | 11.0 | 20 | 12.4* | 12 | 10.0 | 16 | 11.2-11.7* | 26 | 5.8 | 11 | 13.5 | 11 | 8.0 | 25 |
| 8.0* | 20 | 12.4 | 11 | 12.8* | 15 | 12.1 | 10 | 16.7 | 59 | 6.9 | 13 | 16.5 | 52 | 8.6 | 42 |
| 8.6 | 57 | 13.8 | 9 | 17.6 | 20 | 12.8 | 6 | 17.5* | 33 | 7.9 | 53 | 17.6-18.0* | 35 | 8.9 | 25 |
| 9.9 | 22 | 16.8 | 14 | 18.3* | 43 | 13.8 | 4 | 19.3* | 55 | 9.2 | 56 | 19.7 | 82 | 9.9 | 36 |
| 16.6 | 42 | 18.5 | 100 | 19.3 | 100 | 15.1 | 13 | 21.4* | 100 | 9.5 | 50 | 22.3 | 100 | 10.4* | 24 |
| 19.0 | 27 | 19.7* | 22 | 22.2* | 14 | 16.7* | 64 | 22.4* | 33 | 10.3* | 13 | 23.2 | 26 | 12.5 | 18 |
| 21.1 | 35 | 20.9 | 14 | 23.4* | 23 | 18.6* | 100 | 23.2* | 63 | 11.8 | 20 | 24.4 | 28 | 13.9* | 9 |
| | | 25.0* | 15 | 23.8* | 26 | 20.3* | 79 | 29.0* | 15 | 16.1 | 13 | 25.8 | 17 | 16.2 | 10 |
| | | | | 25.5* | 16 | 21.2 | 24 | | | 16.9 | 39 | 26.5 | 30 | 17.8 | 70 |
| | | | | | | 21.9 | 30 | | | 19.1 | 100 | 27.3 | 31 | | |
| | | | | | | 22.4 | 19 | | | | | | | | |

* Broad

^a Relative intensity for Form V 4.9 (broad) 2 θ is 9; Form VI 13.8 2 θ is 9; Form VIII 12.8 2 θ is 6 and 13.8 2 θ is 4; and Form XII 13.9 (broad) 2 θ is 9 and 27.1 2 θ is 8.

-37-

Table 1. Intensities and Peak Locations of All Diffraction Lines With Relative Intensity Greater Than 10%^a Forms V to XIX
(Measured on Shimadzu Diffractometer)

(Page 2 of 3)

| Form V | | Form VI | | Form VII | | Form VIII | | Form IX | | Form X | | Form XI | | Form XII | |
|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|
| 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) | 2 θ | Relative Intensity (>10%) |
| | | | | | | 25.8 | 33 | 19.8 | 71 | 28.7 | 19 | 19.4 | 100 | | |
| | | | | | | 26.5 | 20 | 21.4 | 49 | 29.5 | 12 | 20.8 | 51 | | |
| | | | | | | 27.4* | 38 | 22.3* | 36 | 30.9* | 17 | 21.7 | 13 | | |
| | | | | | | 30.5 | 20 | 23.7* | 37 | 32.8* | 11 | 22.4-22.6* | 18 | | |
| | | | | | | | | 24.4 | 15 | 33.6* | 15 | 24.3 | 19 | | |
| | | | | | | | | 28.7 | 31 | 36.0* | 15 | 25.5 | 24 | | |
| | | | | | | | | | | 38.5* | 14 | 26.2 | 11 | | |
| | | | | | | | | | | | | 27.1 | 8 | | |

* Broad

^a Relative intensity for Form V 4.9 (broad) 2 θ is 9; Form VI 13.8 2 θ is 9; Form VIII 12.8 2 θ is 6 and 13.8 2 θ is 4; and Form XII 13.9 (broad) 2 θ is 9 and 27.1 2 θ is 8.

Table 1. Intensities and Peak Locations of All Diffraction Lines With Relative Intensity Greater Than 10%^a for Forms V to XIX
(Measured on Shimadzu Diffractometer)

| Form XIII | | | Form XIV | | | Form XV | | | Form XVI | | | Form XVII | | | Form XVIII | | | Form XIX | | |
|------------|---------------------------|--|------------|---------------------------|--|------------|---------------------------|--|------------|---------------------------|--|------------|---------------------------|--|------------|---------------------------|--|------------|---------------------------|--|
| 2 θ | Relative Intensity (>10%) | | 2 θ | Relative Intensity (>10%) | | 2 θ | Relative Intensity (>10%) | | 2 θ | Relative Intensity (>10%) | | 2 θ | Relative Intensity (>10%) | | 2 θ | Relative Intensity (>10%) | | 2 θ | Relative Intensity (>10%) | |
| 8.4 | 100 | | 5.4 | 41 | | 5.7 | 26 | | 5.2 | 37 | | 5.0 | 27 | | 8.0 | 100 | | 5.2 | 32 | |
| 8.9 | 82 | | 6.7 | 31 | | 6.1 | 21 | | 6.4 | 34 | | 6.1 | 33 | | 9.2* | 52 | | 6.3 | 28 | |
| 15.7* | 45 | | 7.7 | 100 | | 6.8 | 18 | | 7.5 | 100 | | 7.3 | 100 | | 9.7* | 40 | | 7.0 | 100 | |
| 16.4* | 46 | | 8.1 | 35 | | 7.5 | 39 | | 8.7 | 79 | | 7.9 | 30 | | 12.1 | 24 | | 8.6 | 74 | |
| 17.6* | 57 | | 9.0 | 65 | | 8.1 | 39 | | 10.5* | 19 | | 8.5 | 29 | | 16.6* | 48 | | 10.5 | 34 | |
| 18.1* | 62 | | 16.5* | 15 | | 8.5 | 42 | | 12.0* | 10 | | 9.1 | 22 | | 18.5 | 67 | | 11.6* | 26 | |
| 19.7* | 58 | | 17.6* | 17 | | 9.5 | 33 | | 12.7* | 17 | | 10.0 | 45 | | | | | 12.7* | 35 | |
| 20.8* | 91 | | 18.0-18.7* | 21 | | 10.5* | 18 | | 16.7 | 26 | | 12.1* | 24 | | | | | 14.0 | 15 | |
| 23.8* | 57 | | 19.5* | 18 | | 19.1-19.6* | 32 | | 18.3* | 27 | | 14.8 | 17 | | | | | 16.7* | 30 | |
| | | | | | | | | | 19.5 | 23 | | 16.0-16.5* | 20 | | | | | 18.9 | 86 | |
| | | | | | | | | | 20.1-20.4* | 37 | | 17.5* | 28 | | | | | 20.8 | 94 | |
| | | | | | | | | | 21.2-21.9* | 32 | | 19.0* | 46 | | | | | 23.6* | 38 | |
| | | | | | | | | | 22.9-23.3* | 38 | | 19.5 | 65 | | | | | 25.5* | 32 | |
| | | | | | | | | | 24.4-25.0* | 35 | | 20.2* | 47 | | | | | | | |
| | | | | | | | | | | | | 21.3 | 64 | | | | | | | |
| | | | | | | | | | | | | 21.6 | 55 | | | | | | | |
| | | | | | | | | | | | | 22.0 | 45 | | | | | | | |

* Broad

Forms XIV, XV, XVI, XVII, XVIII, and XIX were measured on Bruker D-5000 Diffractometer.

^a Relative intensity for Form V 4.9 (broad) 2 θ is 9; Form VI 13.8 2 θ is 9; Form VIII 12.8 2 θ is 6 and 13.8 2 θ is 4; and Form XII 13.9 (broad) 2 θ is 9 and 27.1 2 θ is 8.

-39-

Because only 19 crystalline forms of atorvastatin are known, each form can be identified and distinguished from the other crystalline forms by either a combination of lines or a pattern that is different from the X-ray powder diffraction of the other forms.

- 5 For example, Table 2 lists combination of 2 θ peaks for Forms V to XIX atorvastatin, i.e., a set of X-ray diffraction lines that are unique to each form. Forms I to IV atorvastatin disclosed in United States Patent Numbers 5,969,156 and 6,121,461 are included for comparison.

Table 2. Forms I to XIX Unique Combination of 2θ Peaks

| Form I | Form II | Form III | Form IV | Form V | Form VI | Form VII | Form VIII | Form IX | Form X |
|--------|-----------|----------|---------|--------|---------|----------|-----------|---------|--------|
| 9.0 | 8.5 | 8.3 | 4.7 | 6.0 | 7.2 | 8.6 | 7.5 | 8.8 | 4.7 |
| 9.3 | 9.0 | 16.4 | 5.2 | 7.0 | 8.3 | 10.2 | 9.2 | 9.4* | 6.9 |
| 10.1 | 17.1-17.4 | 19.9 | 7.7 | 8.0* | 11.0 | 12.8* | 10.0 | 16.7 | 7.9 |
| 10.4 | 20.5 | 24.2 | 9.4 | 9.9 | 18.5 | 17.6 | 16.7* | 17.5* | 9.2 |
| 11.7 | | | 10.1 | 16.6 | | 18.3* | 18.6* | 19.3* | 9.5 |
| 12.0 | | | | | | 19.3 | 20.3* | 21.4* | 19.1 |
| 16.8 | | | | | | | | 29.0* | 19.8 |
| | | | | | | | | 30.0 | |

-40-

| Form XI | Form XII | Form XIII | Form XIV | Form XV | Form XVI | Form XVII | Form XVIII | Form XIX |
|---------|----------|-----------|----------|------------|------------|-----------|------------|----------|
| 10.8* | 7.7 | 8.4 | 5.4 | 5.7 | 5.2 | 6.1 | 8.0 | 5.5 |
| 16.5 | 8.0 | 8.9 | 6.7 | 6.1 | 6.4 | 7.3 | 9.2* | 7.0 |
| 19.7 | 8.6 | 20.8* | 7.7 | 7.5 | 7.5 | 7.9 | 16.6* | 8.6 |
| 22.3 | 8.9 | 23.8* | 8.1 | 8.1 | 8.7 | 10.0 | 18.5 | 10.5 |
| | 9.9 | | 9.0 | 8.5 | 16.7 | 19.0* | | 12.7* |
| | 17.8 | | | 9.5 | 20.1-20.4* | 19.5 | | 18.9 |
| | 19.4 | | | 19.1-19.6* | 22.9-23.3* | 21.3 | | 20.8 |
| | | | | | | 21.6 | | |

* Broad

Forms I to XIII were measured on Shimadzu XRD-6000 diffractometer. Forms XIV to XIX were measured on Bruker D 5000I diffractometer. Form II 2θ peaks from US Patent Number 5,969,156.


-41-

SOLID STATE NUCLEAR MAGNETIC RESONANCE (NMR)

Methodology

Solid-state ^{13}C NMR spectra were obtained at 270 or 360 MHz Tecmag instruments. High-power proton decoupling and cross-polarization with magic-angle spinning at approximately 4.7 and 4.2 kHz or 4.6 and 4.0 kHz were used for 68 MHz (^{13}C frequency) data acquisition, 4.9 and 4.4 kHz were used for 91 MHz (^{13}C frequency) data acquisition. The magic angle was adjusted using the Br signal of KBr by detecting the side bands. A sample was packed into a 7 mm Doty rotor and used for each experiment. The chemical shifts were referenced externally to adamantane except for Form X where the chemical shifts are arbitrary.

Table 3 shows the solid-state NMR spectrum for crystalline Forms V, VI, VII, VIII, and X atorvastatin.



-42-

Table 3. Chemical Shifts for Forms V, VI, VII, VIII, and X Atorvastatin

| Chemical Shift | | | | |
|----------------|-------|-------|-------|-------|
| V | VI | VII | VIII | X |
| 185.7 | | 186.5 | 186.1 | 187.0 |
| | | 183.3 | 179.5 | |
| 176.8 | 176.5 | 176.8 | | 179.5 |
| 166.9 | 168.2 | 166.5 | 167.9 | 165.5 |
| | 163.1 | | 161.0 | |
| | 159.8 | 159.2 | | 159.4 |
| 138.7 | 136.8 | 137.6 | 139.4 | 137.9 |
| 136.3 | | | 132.9 | 134.8 |
| 133.0 | | | | |
| | | | | 129.4 |
| 128.4 | 127.8 | 128.3 | 128.7 | 127.9 |
| | | | 124.7 | 123.2 |
| 122.0 | 122.3 | 122.3 | | |
| | | | 121.8 | |
| | 118.8 | 119.2 | | 119.9 |
| 117.0 | | | | |
| 116.3 | | | 116.6 | |
| | 113.7 | | | |
| | 88.2 | 74.5 | | |
| | 79.3 | | | |
| | 70.5 | 70.3 | | 71.1 |
| 68.0 | | 68.3 | 67.0 | |
| | | 66.2 | | |
| 43.1 | 43.3 | 43.5 | 43.3 | 43.7 |
| | | 40.3 | | |
| | 36.9 | | | 40.9 |
| | 31.9 | | | |

-43-

Table 3. Chemical Shifts for Forms V, VI, VII, VIII and X Atorvastatin (cont)

| Chemical Shift | | | | |
|----------------|------|------|------|------|
| V | VI | VII | VIII | X |
| 25.6 | 25.9 | 26.3 | 26.7 | 26.4 |
| | | 24.9 | 24.7 | 25.3 |
| | 22.5 | 20.2 | 20.9 | 20.3 |
| 19.9 | | | 20.1 | |
| | | | | 18.3 |

Forms V, VI, VII, VIII, and X: Relative peak intensity over 20 are shown here (4.5, 4.6, 4.7, or 4.9 kHz CPMAS). Spectra were obtained using two different magic-angle spinning rates to determine spinning sidebands.

Form X: Relative peak intensity over 20 are shown here (5.0 kHz CPMAS).

Table 4 shows unique solid-state NMR peaks for Forms V, VI, VII, VIII and X atorvastatin, ie, peaks within ± 1.0 ppm. Forms I to IV atorvastatin are included for comparison.

Table 4. Forms I to VIII and X Unique Solid-State NMR Peaks

| Form I | Form II | Form III | Form IV | Form V | Form VI | Form VII | Form VIII | Form X |
|--------|---------|----------|---------|--------|---------|----------|-----------|--------|
| 182.8 | 181.0 | 161.0 | 181.4 | 176.8 | 163.1 | 183.3 | 132.9 | 18.3 |
| 131.1 | 163.0 | 140.1 | 63.5 | | 36.9 | 176.8 | | |
| 73.1 | 161.0 | 131.8 | 17.9 | | 31.9 | 74.5 | | |
| 64.9 | 140.5 | 69.8 | | | | | | |
| | | 35.4 | | | | | | |

RAMAN SPECTROSCOPY

Methodology

The Raman spectrum was obtained on a Raman accessory interfaced to a Nicolet Magna 860 Fourier transform infrared spectrometer. The accessory
5 utilizes an excitation wavelength of 1064 nm and approximately 0.45 W of neodymium-doped yttrium aluminum garnet (Nd:YAG) laser power. The spectrum represents 64 or 128 co-added scans acquired at 4 cm^{-1} resolution. The sample was prepared for analysis by placing a portion into a 5-mm diameter glass tube and positioning this tube in the spectrometer. The spectrometer was
10 calibrated (wavelength) with sulfur and cyclohexane at the time of use.

Table 5 shows the Raman spectra for Forms V, VI, VII, VIII, X, and XII atorvastatin.

-46-

Table 5. Raman Peak Listing for Forms V, VI, VII, VIII, X and XII
Atorvastatin

| Form V | Form VI | Form VII | Form VIII | Form X | Form XII |
|--------|---------|----------|-----------|--------|----------|
| 3062 | 3058 | 3060 | 3065 | 3062 | 3064 |
| | | | | | 2973 |
| | 2935 | 2927 | 2923 | 2911 | 2926 |
| 1652 | 1651 | 1649 | 1658 | 1650 | 1652 |
| 1604 | 1603 | 1603 | 1603 | 1603 | 1603 |
| 1528 | 1556 | 1524 | 1531 | 1525 | 1527 |
| | 1525 | | 1510 | | |
| | | | 1481 | | |
| 1478 | 1478 | 1476 | | 1478 | 1470 |
| 1440 | | | | | |
| 1413 | 1412 | 1412 | 1413 | 1411 | 1410 |
| 1397 | | 1397 | | | |
| 1368 | | 1368 | | 1369 | 1367 |
| | | | | 1240 | 1240 |
| 1158 | 1157 | 1159 | | 1158 | 1159 |
| 1034 | | 1034 | | 1034 | 1034 |
| 1001 | 997 | 998 | 997 | 999 | 1002 |
| 825 | | 824 | | 824 | 823 |
| 245 | | | | | |
| 224 | | | | | |
| 130 | | | | | |
| | | 114 | 121 | 116 | |

Relative peak intensity over 20 are shown.

Table 6 lists unique Raman peaks for Forms V, VI, VII, VIII, X, and XII atorvastatin, ie, only one other form has a peak with $\pm 4 \text{ cm}^{-1}$. In the case of Forms VI and X, it is a unique combination of peaks. Forms I to IV atorvastatin are included for comparison.

Table 6. Forms I to VIII, X and XII Unique Raman Peaks

| Form I | Form II | Form III | Form IV | Form V | Form VI* | Form VII | Form VIII | Form X* | Form XII |
|--------|---------|----------|---------|--------|----------|----------|-----------|---------|----------|
| 3080 | 1663 | 2938 | 423 | 1440 | 3058 | 1397 | 1510 | 3062 | 2973 |
| 1512 | 359 | 1660 | 215 | 1397 | 2935 | | 1481 | 2911 | |
| 1439 | | 1510 | 132 | 130 | 1556 | | 1413 | 1525 | |
| 142 | | 1481 | | | 1525 | | 121 | 1240 | |
| | | 1427 | | | | | | | |
| | | 1182 | | | | | | | |
| | | 859 | | | | | | | |

* Unique combination of Raman peaks

-48-

Crystalline Forms V to XIX atorvastatin of the present invention may exist in anhydrous forms as well as hydrated and solvated forms. In general, the hydrated forms are equivalent to unhydrated forms and are intended to be encompassed within the scope of the present invention. Crystalline Form XIV contains about 6 mol of water. Preferably, Form XIV contains 6 mol of water. Crystalline Forms V, X, and XV atorvastatin contain about 3 mol of water. Preferably, Forms V, X, and XV atorvastatin contain 3 mol of water.

Crystalline Form VII contains about 1.5 mol of water. Preferably, Form VII atorvastatin contains 1.5 mol of water. Crystalline Form VIII contains about 2 mol of water. Preferably, Form VIII atorvastatin contains 2 mol of water.

Crystalline Forms XVI-XIX may exist as a solvate.

Crystalline forms of atorvastatin of the present invention, regardless of the extent of hydration and/or solvation having equivalent x-ray powder diffractograms, ssNMR, or Raman spectra are within the scope of the present invention.

Crystalline forms, in general, can have advantageous properties. A polymorph, solvate, or hydrate is defined by its crystal structure and properties. The crystal structure can be obtained from X-ray data or approximated from other data. The properties are determined by testing. The chemical formula and chemical structure does not describe or suggest the crystal structure of any particular polymorphic or crystalline hydrate form. One cannot ascertain any particular crystalline form from the chemical formula, nor does the chemical formula tell one how to identify any particular crystalline solid form or describe its properties. Whereas a chemical compound can exist in three states—solid, solution, and gas-crystalline solid forms exist only in the solid state. Once a chemical compound is dissolved or melted, the crystalline solid form is destroyed and no longer exists (Wells J.I., Aulton M.E. *Pharmaceutics. The 'Science of Dosage Form Design. Reformulation*, Aulton M.E. ed., Churchill Livingstone, 1988;13:237).

The new crystalline forms of atorvastatin described herein have advantageous properties. Form VII has good chemical stability, which is comparable to Form I (disclosed in United States Patent Number 5,969,156).

-49-

Since noncrystalline forms of atorvastatin are not chemically stable, this is a significant advantage, which would translate into enhanced shelf life and longer expiration dating. Form VII can be prepared from acetone/water, whereas Form I is prepared from the more toxic methanol/water system. Form VII is the
5 sesquihydrate and contains less water, meaning that a unit weight of Form VII contains more atorvastatin molecules, meaning it is of higher potency.

The ability of a material to form good tablets at commercial scale depends upon a variety of drug physical properties, such as the Tableting Indices described in Hiestand H. and Smith D., Indices of Tableting Performance, *Powder*
10 *Technology*, 1984;38:145-159. These indices may be used to identify forms of atorvastatin calcium which have superior tableting performance. One such index is the Brittle Fracture Index (BFI), which reflects brittleness, and ranges from 0 (good - low brittleness) to 1 (poor - high brittleness). For example, Form VII has a BFI value 0.09, while Form I has a BFI value 0.81. Thus, Form VII is less brittle
15 than Form I. This lower brittleness indicates greater ease of manufacture of tablets.+

Form VIII also has less water than Form I (dihydrate vs trihydrate) and thus a gram of Form VIII contains more atorvastatin molecules.

Form X is advantageous in that it can be prepared from the less toxic
20 isopropanol (IPA):water system, thus avoiding the more toxic methanol:water system.

Form XII has the highest melting point (210.6). Since high melting point correlates with stability at high temperature, this means this form is most stable at temperatures near the melting point. High melting forms can be advantageous
25 when process methods involving high temperatures are used. Form XII is also prepared from the less toxic tetrahydrofuran (THF) water system.

Form XIV is prepared using the less toxic THF/water system.

The present invention provides a process for the preparation of crystalline Forms V to XIX atorvastatin which comprises crystallizing atorvastatin from a
30 solution in solvents under conditions which yield crystalline Forms V to XIX atorvastatin.

-50-

The precise conditions under which crystalline Forms V to XIX atorvastatin are formed may be empirically determined, and it is only possible to give a number of methods which have been found to be suitable in practice.

5 The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the
10 compounds of the present invention can be administered transdermally. It will be obvious to those skilled in the art that the following dosage forms may comprise as the active component, either compounds or a corresponding pharmaceutically acceptable salt of a compound of the present invention.

For preparing pharmaceutical compositions from the compounds of the
15 present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents,
20 or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape
25 and size desired.

The powders and tablets preferably contain from two or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax,
30 cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component, with or without other carriers,

-51-

is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

5 For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

10 Liquid form preparations include solutions, suspensions, retention enemas, and emulsions, for example water or water propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

15 Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

20 Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, 25 solubilizing agents, and the like.

The pharmaceutical preparation is preferably in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as 30 packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

-52-

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.5 mg to 100 mg, preferably 2.5 mg to 80 mg according to the particular application and the potency of the active component. The composition can, if desired, also contain other compatible therapeutic agents.

5 In therapeutic use as hypolipidemic and/or hypocholesterolemic agents and agents to treat osteoporosis and Alzheimer's disease, the crystalline Forms V to XIX atorvastatin utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 2.5 mg to about 80 mg daily. A daily dose range of about 2.5 mg to about 20 mg is preferred. The dosages, however, 10 may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the 15 optimum effect under the circumstance is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The following nonlimiting examples illustrate the inventors' preferred methods for preparing the compounds of the invention.

EXAMPLE 1

20 [R-(R*,R*)]-2-(4-Fluorophenyl)- β , δ -dihydroxy-5-(1-methylethyl)-3-phenyl-4-
[(phenylamino)carbonyl]-1H-pyrrole-1-heptanoic acid hemi calcium salt
(Forms V-XIX atorvastatin)

Form V Atorvastatin

Method A

25 Amorphous atorvastatin calcium (United States Patent Number 5,273,995) was slurried in a mixture of acetonitrile/water (9:1) to afford crystalline Form V atorvastatin.

-53-

Method B

Crystalline Form I atorvastatin calcium (United States Patent Number 5,969,156) was slurried in a mixture of acetonitrile/water (9:1) at 60°C overnight, filtered, and air dried to afford crystalline Form V atorvastatin.

5 Method C

Amorphous atorvastatin calcium (United States Patent Number 5,273,995) was stressed under vapors of acetonitrile/water (9:1) to afford crystalline Form V atorvastatin.

Method D

10 Acetonitrile was added to a solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in tetrahydrofuran/water (9:1) and cooled to afford crystalline Form V atorvastatin.

Method E

15 Acetonitrile was added to a solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in dimethylformamide/water and fast evaporation affords crystalline Form V atorvastatin.

Method F

20 Amorphous atorvastatin calcium (United States Patent Number 5,273,995) diffused in a vapor of acetonitrile/water (9:1) to afford crystalline Form V atorvastatin.

Crystalline Form V atorvastatin, mp 171.4°C, trihydrate
Karl Fischer 4.88% (3 mol of water).

Form VI AtorvastatinMethod A

25 Amorphous atorvastatin calcium (United States Patent Number 5,273,995) was placed into a vapor jar containing dimethylformamide/water (9:1) for 20 days to afford crystalline Form VI atorvastatin.

-54-

Method B

Fast evaporation of a dimethylformamide/water solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) afforded crystalline Form VI atorvastatin.

5 Method C

Fast evaporation of a dimethylformamide/water (saturated) solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) seeded with crystalline Form VI afforded crystalline Form VI atorvastatin.

Crystalline Form VI atorvastatin, mp 145.9°C.

10 **Form VII Atorvastatin**Method A

A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (1:1) (5.8 mg/mL) was stirred overnight. A solid formed which was filtered to afford crystalline Form VII atorvastatin.

15 Method B

A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (1:1) was evaporated at 50°C to afford crystalline Form VII atorvastatin.

Method C

20 A saturated solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (1:1) was seeded with crystalline Form VII atorvastatin to afford crystalline Form VII atorvastatin.

Method D

25 Fast evaporation of a saturated solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (1:1) was seeded with crystalline Form VII to afford crystalline Form VII atorvastatin.

Crystalline Form VII atorvastatin, mp 195.9°C, 1.5 hydrate
Karl Fischer 2.34% (1.5 mol of water).

-55-

Form VIII AtorvastatinMethod A

5 A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in dimethylformamide/water (saturated) (9:1), was seeded with crystalline Form VII and evaporated to afford crystalline Form VIII atorvastatin.

Method B

10 Fast evaporation of a solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in dimethylformamide/water (9:1) affords crystalline Form VIII atorvastatin.

Crystalline Form VIII atorvastatin, mp 151°C, dihydrate
Karl Fischer 2.98% (2 mol of water).

Form IX AtorvastatinMethod A

15 A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (6:4) (3.4 mg/mL) was evaporated on a rotary evaporator to afford crystalline Form IX atorvastatin.

Method B

20 A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (6:4) was filtered, seeded with crystalline Form IX evaporated on a rotary evaporator to afford crystalline Form IX atorvastatin.

Method C

25 A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetone/water (6:4) was stirred for 0.5 hours, filtered, evaporated on rotary evaporator to concentrate the solution, and dried in a vacuum oven to afford crystalline Form IX atorvastatin.

-56-

Form X AtorvastatinMethod A

5 A slurry of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in isopropanol/water (9:1) was stirred for a few days, filtered, and air dried to afford crystalline Form X atorvastatin.

Method B

A slurry of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in isopropanol/water (9:1) was stirred for 5 days, filtered, and air dried to afford crystalline Form X atorvastatin.

10 Method C

A saturated solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in isopropanol/water (9:1) was stirred for 2 days, filtered, and air dried to afford crystalline Form X atorvastatin.

15 Crystalline Form X atorvastatin, mp 180.1°C, trihydrate
Karl Fischer 5.5% (3.5 mol of water).

Form XI Atorvastatin

A solution of amorphous atorvastatin calcium (United States Patent Number 5,273,995) in acetonitrile/water (9:1) was filtered and allowed to evaporate slowly to afford crystalline Form XI atorvastatin.

20 **Form XII Atorvastatin**

Crystalline Form I atorvastatin calcium (United States Patent Number 5,969,156) was slurried in tetrahydrofuran/water (2:8) at 90°C for 5 days, filtered, and air dried to afford crystalline Form XII atorvastatin.

Crystalline Form XII atorvastatin, mp 210.6°C.

25 **Form XIII Atorvastatin**

Crystalline Form I atorvastatin calcium (United States Patent Number 5,969,156) was added to 10 mL 2:8 water:methanol to leave a layer of solid on the bottom of a vial. The slurry was heated to about 70°C for 5 days. The supernatant was removed, and the solid air dried to afford crystalline Form XIII atorvastatin.

-57-

Form XIV Atorvastatin

Amorphous atorvastatin calcium (United States Patent Number 5,273,995), 1 g, was slurried for 3 weeks in 45 mL of isopropyl alcohol/5 mL of water (9:1) at ambient temperature. The mixture was filtered to afford crystalline Form XIV atorvastatin after drying at ambient temperature.

Differential scanning calorimetry (DSC) indicates a low desolvation event at about 60°C (peak) followed by a melt at about 150°C. Combustion analysis indicates that the compound is a hexahydrate. Thermographic infrared spectroscopy (TG-1R) shows the compound contains water. Karl Fischer shows the compound contains 5.8% water.

Form XV Atorvastatin

Amorphous atorvastatin calcium (United States Patent Number 5,273,995), 1 g, was slurried for 3 weeks in 45 mL acetonitrile/5 mL of water (9:1) at ambient temperature. The mixture was filtered to afford crystalline Form XV atorvastatin after drying at ambient temperature. DSC indicates a low desolvation event at about 78°C (peak) followed by a melt at about 165°C. Combustion analysis indicates that the compound is a trihydrate. TG-1R shows the compound contains water.

Form XVI Atorvastatin

Amorphous atorvastatin calcium (United States Patent Number 5,273,995), 1 g, was slurried for about 1 day in 9:1 acetonitrile/water at room temperature. The mixture was filtered to afford crystalline Form XVI atorvastatin after drying at ambient temperature. DSC indicates a broad endotherm at peak temperature of 72°C and an endotherm with onset temperature of 164°C. The weight loss profile by thermographic analysis (TGA) indicates a total weight loss of about 7% at 30°C to 160°C. Combustion analysis indicates that TGA and Karl Fischer analysis (shows 7.1% water) indicates the compound is a tetrahydrate/acetonitrile solvate.

Form XVII Atorvastatin

Amorphous atorvastatin calcium (United States Patent Number 5,273,995), 0.5 g, was slurried for about 2 days in 5 mL of 9:1 dimethylformamide (DMF)/water containing 25 mL of acetonitrile at room temperature. The mixture

-58-

was filtered to afford crystalline Form XVII atorvastatin after drying at ambient temperature. DSC showed multiple broad endotherms indicating the compound was a solvate.

Form XVIII Atorvastatin

- 5 Crystalline Form XVI atorvastatin, 0.5 g, was dried for about 1 day at room temperature to afford crystalline Form XVIII atorvastatin. DSC showed a broad endotherm at low temperature indicating the compound was a solvate. Karl Fischer analysis showed the compound contained 4.4% water.

Form XIX Atorvastatin

- 10 Amorphous atorvastatin calcium (United States Patent Number 5,273,995), 0.4 g, was slurried for about 7 days in 4 mL methyl ethyl ketone at room temperature. The mixture was filtered to afford crystalline Form XIX atorvastatin after drying at ambient temperature. DSC indicated a low desolvation event at about 50°C (peak) followed by a melt at about 125°C. TGA analysis indicates that
- 15 the compound is a solvate that desolvates at low temperature.

-59-

CLAIMS

What is claimed is:

1. A crystalline Form V atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 4.9 (broad), 6.0, 7.0, 8.0 (broad), 8.6, 9.9, 16.6, 19.0, and 21.1.
2. A crystalline Form VI atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 7.2, 8.3, 11.0, 12.4, 13.8, 16.8, 18.5, 19.7 (broad), 20.9, and 25.0.
3. A crystalline Form VII atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 8.6, 10.2, 12.4 (broad), 12.8 (broad), 17.6, 18.3 (broad), 19.3, 22.2 (broad), 23.4 (broad), 23.8 (broad), and 25.5 (broad).
4. A crystalline Form VIII atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 7.5, 9.2, 10.0, 12.1, 12.8, 13.8, 15.1, 16.7 (broad), 18.6 (broad), 20.3 (broad), 21.2, 21.9, 22.4, 25.8, 26.5, 27.4 (broad), and 30.5.
5. A crystalline Form IX atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 8.8, 9.4 (broad), 11.2-11.7 (broad), 16.7, 17.5 (broad), 19.3 (broad), 21.4 (broad), 22.4 (broad), 23.2 (broad), and 29.0 (broad).
6. A crystalline Form X atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using

-60-


CuK α radiation: 4.7, 5.2, 5.8, 6.9, 7.9, 9.2, 9.5, 10.3 (broad), 11.8, 16.1, 16.9, 19.1, 19.8, 21.4, 22.3 (broad), 23.7 (broad), 24.4, and 28.7.

- 5 7. A crystalline Form XI atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 10.8 (broad), 12.0, 13.5, 16.5, 17.6-18.0 (broad), 19.7, 22.3, 23.2, 24.4, 25.8, 26.5, 27.3, 28.7, 29.5, 30.9 (broad), 32.8 (broad), 33.6 (broad), 36.0 (broad), and 38.5 (broad).
- 10 8. A crystalline Form XII atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 5.4, 7.7, 8.0, 8.6, 8.9, 9.9, 10.4 (broad), 12.5, 13.9 (broad), 16.2, 17.8, 19.4, 20.8, 21.7, 22.4-22.6 (broad), 24.3, 25.5, 26.2, and 27.1.
- 15 9. A crystalline Form XIII atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 8.4, 8.9, 15.7 (broad), 16.4 (broad), 17.6 (broad), 18.1 (broad), 19.7 (broad), 20.8 (broad), and 23.8 (broad).
- 20 10. A crystalline Form XIV atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 5.4, 6.7, 7.7, 8.1, 9.0, 16.5 (broad), 17.6 (broad), 18.0-18.7 (broad), and 19.5 (broad).
11. A crystalline Form XV atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using CuK α radiation: 5.7, 6.1, 6.8, 7.5, 8.1, 8.5, 9.5, 10.5 (broad), and 19.1-19.6 (broad).

-61-


12. A crystalline Form XVI atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using $\text{CuK}\alpha$ radiation: 5.2, 6.4, 7.5, 8.7, 10.5 (broad), 12.0 (broad), 12.7 (broad), 16.7, 18.3 (broad), 19.5, 20.1-20.4 (broad), 21.2-21.9 (broad), 22.9-23.3 (broad), and 24.4-25.0 (broad).
13. A crystalline Form XVII atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using $\text{CuK}\alpha$ radiation: 5.0, 6.1, 7.3, 7.9, 8.5, 9.1, 10.0, 12.1 (broad), 14.8, 16.0-16.5 (broad), 17.5 (broad), 19.0 (broad), 19.5, 20.2 (broad), 21.3, 21.6, and 22.0.
14. A crystalline Form XVIII atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using $\text{CuK}\alpha$ radiation: 8.0, 9.2 (broad), 9.7 (broad), 12.1, 16.6 (broad), and 18.5.
15. A crystalline Form XIX atorvastatin or a hydrate thereof having an X-ray powder diffraction containing the following 2θ values measured using $\text{CuK}\alpha$ radiation: 5.2, 6.3, 7.0, 8.6, 10.5, 11.6 (broad), 12.7 (broad), 14.0, 16.7 (broad), 18.9, 20.8, 23.6 (broad), and 25.5 (broad).

1/35


FIG. 1

2/35


FIG. 2




3/35

FIG. 3

4/35


FIG. 4

5/35

FIG. 5


6/35

FIG. 6




7/35

FIG. 7




8/35

FIG. 8




9/35


FIG. 9

10/35

FIG. 10




11/35

FIG. 11


12/35

FIG. 12




13/35

FIG. 13




14/35

FIG. 14




15/35


FIG. 15




16/35

FIG. 16


17/35

FIG. 17


18/35

FIG. 18


19/35

FIG. 19

20/35


FIG. 20

21/35


FIG. 21

22/35


FIG. 22



23/35


FIG. 23

24/35

FIG. 24


25/35

FIG. 25




26/35

FIG. 26




27/35

FIG. 27




28/35

FIG. 28



29/35

FIG. 29



30/35

FIG. 30






FIG. 31




32/35

FIG. 32



33/35

FIG. 33



34/35

FIG. 34





FIG. 35



INTERNATIONAL SEARCH REPORT

Intern Application No
PCT/IB 02/01796

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D207/34 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| A | WO 97 03959 A (WARNER LAMBERT CO ;BRIGGS CHRISTOPHER A (US); JENNINGS REX ALLEN () 6 February 1997 (1997-02-06) the whole document --- | 1-15 |
| A | WO 97 03958 A (WARNER LAMBERT CO ;MCKENZIE ANN T (US)) 6 February 1997 (1997-02-06) the whole document --- | 1-15 |
| A | WO 01 44181 A (TULLY WILLIAM ;CONNELL JOHN O (IE); MADIGAN EVELYN (IE); WARNER LA) 21 June 2001 (2001-06-21) the whole document --- | 1-15 |
| A | WO 01 44180 A (TULLY WILLIAM ;WARNER LAMBERT RES AND DEV IRE (IE)) 21 June 2001 (2001-06-21) the whole document --- | 1-15 |
| -/-- | | |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

2 July 2002

Date of mailing of the international search report

12/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Schmid, J-C

INTERNATIONAL SEARCH REPORT

Interr. Application No
PCT/IB 02/01796

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| A | WO 01 36384 A (TEVA PHARMA ;AYALON ARI (IL); NIDDAM VALERIE (IL); ROYTBAT SOFIA) 25 May 2001 (2001-05-25) cited in the application the whole document ---- | 1-15 |
| E | WO 02 41834 A (TEVA PHARMA ;TEVA PHARMACEUTICALS USA INC (US)) 30 May 2002 (2002-05-30) the whole document ---- | 1-15 |
| E | WO 02 43732 A (ISHAI ETI ;SAMBURSKY GUY (IL); TEVA PHARMA (IL); ARONHIME JUDITH () 6 June 2002 (2002-06-06) the whole document ----- | 1-15 |

INTERNATIONAL SEARCH REPORT

 Intern: I Application No
 PCT/IB 02/01796

| Patent document cited in search report | | Publication date | Patent family member(s) | Publication date |
|---|---|---------------------|--|--|
| WO 9703959 | A | 06-02-1997 | AT 208375 T AU 725424 B2 AU 6484296 A BG 102187 A BR 9609872 A CA 2220018 A1 CN 1190955 A CZ 9800121 A3 DE 69616808 D1 DE 69616808 T2 DK 848705 T3 EE 9800015 A EP 1148049 A1 EP 0848705 A1 ES 2167587 T3 HR 960339 A1 HU 9900678 A2 IL 122118 A JP 11509230 T NO 980207 A NZ 312907 A PL 324496 A1 PT 848705 T SI 848705 T1 SK 6298 A3 WO 9703959 A1 US 5969156 A | 15-11-2001 12-10-2000 18-02-1997 30-10-1998 23-03-1999 06-02-1997 19-08-1998 14-10-1998 13-12-2001 29-05-2002 04-02-2002 17-08-1998 24-10-2001 24-06-1998 16-05-2002 30-04-1998 28-07-1999 14-07-1999 17-08-1999 16-01-1998 22-12-2000 25-05-1998 28-02-2002 30-04-2002 07-10-1998 06-02-1997 19-10-1999 |
| WO 9703958 | A | 06-02-1997 | AT 207465 T AU 725368 B2 AU 6484196 A BG 102186 A BR 9610567 A CA 2220458 A1 CN 1190957 A , B CZ 9800123 A3 DE 69616358 D1 DK 848704 T3 EE 9800016 A EP 0848704 A1 ES 2166456 T3 HR 960313 A1 HU 9901687 A2 IL 122162 A JP 11509229 T NO 980208 A NZ 312906 A PL 324532 A1 SK 5998 A3 TW 401399 B WO 9703958 A1 US 6121461 A | 15-11-2001 12-10-2000 18-02-1997 30-10-1998 06-07-1999 06-02-1997 19-08-1998 17-06-1998 29-11-2001 04-02-2002 17-08-1998 24-06-1998 16-04-2002 30-04-1998 28-10-1999 14-07-1999 17-08-1999 16-01-1998 22-12-2000 08-06-1998 06-05-1998 11-08-2000 06-02-1997 19-09-2000 |
| WO 0144181 | A | 21-06-2001 | AU 2214301 A WO 0144181 A1 | 25-06-2001 21-06-2001 |
| WO 0144180 | A | 21-06-2001 | AU 2544001 A WO 0144180 A1 | 25-06-2001 21-06-2001 |

INTERNATIONAL SEARCH REPORT

Intern I Application No

PCT/IB 02/01796

| Patent document cited in search report | | Publication date | | Patent family member(s) | Publication date |
|---|---|---------------------|----|----------------------------|---------------------|
| WO 0136384 | A | 25-05-2001 | AU | 1617301 A | 30-05-2001 |
| | | | WO | 0136384 A1 | 25-05-2001 |
| WO 0241834 | A | 30-05-2002 | WO | 0241834 A2 | 30-05-2002 |
| WO 0243732 | A | 06-06-2002 | WO | 0243732 A1 | 06-06-2002 |
| | | | WO | 0243667 A2 | 06-06-2002 |